Resistance to Fusarium Wilt in Common Bean Germplasm. (C01-brick161228-Poster)

Authors:

- M.A.Brick Colorado State University
- J.B.Ogg Colorado State University
- H.F.Schwartz Colorado State University
- P.F.Byrne Colorado State University

Abstract:

Fusarium wilt of dry bean caused by Fusarium oxysporum. f. sp. phaseoli (Fop) has recently become a serious disease on common bean (Phaseolus vulgaris L.) in the High Plains and western U.S. Race variability of Fop exists and host resistance genes differ in their reaction to the different Fop races. Common bean germplasm representative of races from Middle American Center of Diversity was screened for resistance and crosses made to determine modes of inheritance among different resistance genes. Different resistant genes showed allelic differences, and some resistance genes confer resistance to more than one race of Fop. We will identify sources of resistance and modes of inheritance for specific resistance genes.

Corresponding Author Information:

Mark Brick phone: 970-491-6551 Colorado State University fax: 970-491-0564

Department of Soil and Crop e-mail:

Sciences mbrick@lamar.colostate.edu

Fort Collins, CO 80523

Presentation Information:

Presentation Date: Tuesday, November 12, 2002

Presentation Time: 4:00-6:00 pm

Poster Board Number: 1009

Keywords:

dry beans, breeding, disease resistance