Predicting the Effects of Agricultural Management Practices on Subsurface Drainage :Nitrate. (S01ma105428-Oral)

Authors:

- L.Ma* USDA-ARS, Fort Collins, CO
- R.W.Malone USDA-ARS, Ames, IA
- L.R.Ahuja USDA-ARS, Fort Collins, CO
- R.S.Kanwar Iowa State University, Ames, IA

Abstract:

Much of the nitrate in surface waters within the Midwest Cornbelt is from subsurface field drainage. Simulation models are tools that can estimate nitrate reduction from BMP's implemented on tile-drained fields. Quantifying nitrate reduction using models may accelerate adoption of BMP's and deepen our understanding of the processes driving water quality improvements. A first step toward using models for these purposes is to assess the model and proposed agricultural systems using a long-term and complete data set. This presentation investigates nitrate transport from the Nashua, IA research farm using the Root Zone Water Quality Model (RZWQM). Observed data included 10 years of tile flow, chemical concentrations in soil and tile flow, and crop growth from 36 one-acre plots under several different agricultural practices and soil types. Modeling results indicate that RZWQM simulates variation in nitrate transport well compared to observed data, and shows promise in quantifying nitrate in tile flow under different climate (year to year), soil, and management conditions.

Corresponding Author Information:

Liwang Ma USDA-ARS 301 S. Howes, P. O. Box E Fort Collins phone: 970-490-8339 e-mail: ma@gpsr.colostate.edu

Presentation Information:

Presentation Date: Thursday, November 14, 2002 Presentation Time: 9:45 am

Keywords:

Tile flow, RZWQM, Nitrate modeling, Agricultural Management