

# Soil Phosphorus Spatial Distribution in Pastures Receiving Poultry Litter Applications. (S11-washburn152224-Poster)

## Authors:

- C.S.M.Washburn\* - *University of Missouri*
- P.P.Motavalli - *University of Missouri*
- N.Kitchen - *USDA, Columbia, MO*
- D.K.Potter - *USDA-NRCS, Columbia, MO*

## Abstract:

Environmentally-based P management strategies could be improved by delineating management zones incorporating the effects of landscape position on soil morphology, hydrology, and soil P distribution. Three farm pasture sites in SW Missouri receiving long-term poultry litter applications were sampled by landscape position (summit, shoulder, upper backslope, middle backslope, lower backslope, footslope, drain) for soil P (Bray 1 P, CaCl<sub>2</sub> P, total P) at two depths (0-5 cm, 5-15 cm). Low landscape positions (drain, footslope) at Sites 1 and 2 had significantly ( $p < 0.05$ ) lower P concentrations compared to high positions (summit, shoulder, upper backslope, middle backslope, lower backslope) for both soil depths. Conversely, Site 3 had higher soil P concentrations in lower landscape positions (fotslope, lower backslope, middle backslope) compared to higher positions (upper backslope, shoulder, summit), although not always significant for each position at each depth. Differences in P distribution among sites may be due to differences in litter management practices and the significantly higher soil P and more dissected, complex topography at Site 3 compared to Sites 1 and 2.

## Corresponding Author Information:

Carol Washburn  
University of Missouri  
302 ABNR  
Columbia, MO 65211

phone: 573-882-4095  
fax: 573-884-5070  
e-mail: washburnc@missouri.edu

## Presentation Information:

Presentation Date: Tuesday, November 12, 2002

Presentation Time: 10:00 am-12:00 pm

Poster Board Number: 2123

**Keywords:**

soil phosphorus, poultry litter, environmental management